

## Bellwork

Simplify each of the following radicals:

1.  $\sqrt{375}$

2.  $\sqrt{28}$

3.  $\sqrt{1100}$

## Solving Quadratic Equations

Given:

You will be given an expression that will contain a squared variable.

*This will be the highest power given.*

Looking for:

The zeros, or roots, of the function.

These are also known as the x-intercepts

## **Processes in Solving Quadratics**

There are several approaches to solving quadratics including:

1. Solving using Square Roots

**This will be the focus of this lesson.**

2. Solving by Factoring

3. Solving by Completing the Square

4. Solving by Using the Quadratic Formula

## **Square Roots Fact**

Did you know there is technically 2 answers when you take the square root of a number?

1. The Principle Root:

This is the root that you are use to.  
The positive root.

2. The Negative Root:

If you square a negative number you get a positive number right?

## Solving by Square Roots

Procedure:

1. Isolate the term that is squared.

[NOTE: This could be a term or an expression!]

2. Take the square root of both sides. Remember that this will now become + or - to give us the principle and the negative roots.

[Simplify the radical if possible.]

3. If there is a square root in the denominator rationalize it by multiplying the top and bottom by the square root that is left.

4. Solve for x.

[This may including having to add/subtract a number that is subtracted/added to the x term, or multiplying/dividing a number that x is divided by or multiplied by.]

5. Simplify if possible

## Example

Solve the following by using square roots:

1.  $12x^2 = 1728$

2.  $7x^2 + 5 = 15$

## Example

Solve the following by using square roots:

3.  $2x^2 + 9 = 105$

4.  $3(x - 4)^2 = 432$

## Example

Solve the following by using square roots:

5.  $2(x + 8)^2 + 6 = -36$